



## Evaluation of the effect of chain training methods on improvement of household awareness and attitude toward healthy nutrition

Farzam Bidarpoor<sup>1</sup>, Soraya Amani<sup>1</sup>, Fatemeh Darabi<sup>2</sup>

1 Deputy of Health, Kurdistan University of Medical Sciences, Sanandaj, Iran

3 PhD Candidate, Department of Health Education, School of Health, Gonabad University of Medical Sciences, Gonabad, Iran

### Original Article

#### Abstract

**BACKGROUND:** Cardiovascular diseases (CVD) are one of the major causative factors of morbidity and mortality in the world which can be easily prevented through lifestyle changes. The aim of this study was to evaluate the effect of chain training methods on increasing household awareness on and attitude toward healthy nutrition in order to prevent CVD.

**METHODS:** This study was performed as a community trial. For the purpose of this study, 4 areas were selected and 4 educational methods were conducted in these areas. The participants were told to share their information with their neighbors and relatives in that area (chain training). Primary knowledge of the residents of these areas before and after the intervention were evaluated in a randomly selected sample (n = 400). Data were analyzed using independent t-test, chi-square, and Fisher's exact test.

**RESULTS:** Pre-intervention and post-intervention mean of knowledge were  $16.42 \pm 3.5$  and  $16.4 \pm 3.6$ , respectively (P = 0.025). Pre-intervention and post-intervention mean of knowledge did not differ in area 1, 2, and 4. A statistically significant increase in knowledge was only observed in area 3 (P < 0.001).

**CONCLUSION:** The use of chain training method by non-governmental organization (NGOs) may not be effective in increasing awareness on healthy lifestyle. Therefore, training through public classes and direct education by health personnel is recommended.

**KEYWORDS:** Training, Knowledge, Behavior, Nutrition

**Date of submission:** 18 Nov 2013, **Date of acceptance:** 5 Jan 2015

**Citation:** Bidarpoor F, Darabi F, Amani S. Evaluation of the effect of chain training methods on improvement of household awareness on and attitude toward healthy nutrition. Chron Dis J 2015; 3(1): 1-6.

### Introduction

Cardiovascular diseases (CVD) are one of the major factors of illness and mortality in the world.<sup>1,2</sup> The most important causative factors of CVD include unhealthy diet, physical inactivity, and tobacco consumption, which can be prevented through appropriate training.<sup>3,4</sup> Recently, lifestyle changes have resulted in some nutritional risk factors in different communities such as Iran.<sup>5-8</sup> Therefore, the general guideline of the American Heart Association has emphasized

the improvement of regional policies and community training in order to change their lifestyles as the main components of CVD prevention.<sup>9,10</sup>

The different methods of health education include lectures, small group discussions, role playing, performing, film, direct observation, face-to-face training, procedures, and etcetera. One of these methods is chain training. In this method, a group are trained and they will transmit the educational materials to others<sup>11</sup>. Today, some political issues and news spread over a population through this method. Now, the question is: "Can chain method be useful in health

#### Corresponding Author:

Fatemeh Darabi

Email: fatemedarabi43@yahoo.com

education?". Answering this question can be effective in teaching of health issues. Nevertheless, since there are not enough personnel, educating the society members by the health system is quite difficult. Evidently, this problem can be solved by society members' training others through chain education method.<sup>12</sup> The aim of this study was to investigate the effect of some training methods on knowledge promotion through chain education method.

### Materials and Methods

The present study was performed as a community trial in the two cities of Sanandaj and Saqez, Kurdistan Province, Western Iran, in 2012 after being approved by the ethics committee of Kurdistan University of Medical Sciences.

In these two cities, four areas were non-

randomly selected based on the experts' opinions. The study was conducted in cooperation with four main non-governmental organizations (NGOs), which were experienced in health planning, and in four stages. First, for the uniformity of the education in all the intervention groups, the educational texts were prepared by nutrition experts of the health system in order to be used by the trainer. In the second stage, the primary knowledge status of the people before the intervention was evaluated in a 400-subject sample size randomly selected based on postal code from all areas. In the third stage, training and intervention programs were conducted by the selected NGOs which had previously participated in similar tasks based on the educational methods listed in table 1.

**Table 1. List of intervention field in the study**

City	The name of the area	Intervention methods	Teaching methods	Type of trainer	Participants in pre-/post-intervention
Sanandaj	Abbasabad	Providing 500 households with face-to-face training and educational CDs with an instruction card	Based on need assessment	Company working in educational and social interface	100-100
Sanandaj	Kanikozaleh	Training parents in schools (each individual only once)	Community-based education or local groups (CBO)	Social interface	100-100
Sanandaj	Farajah	Training through public classes and practical work in the review of food units consumed in the previous few days, measuring anthropometric indices, informing individuals about them, and practical training on cooking	Training in public classes and practical work	Health Personnel	50-50
Saqez	Shakarriz	Training of 323 households through public classes and private sessions, re-training by presenting weekly paper (24 cards per week), and practical nutrition education classes	Leitner method	Health interface	150-150

The participants were also told to share their information with their neighbors and relatives in that area (chain training). The standard questionnaire of the Heart Research Center of Esfahan<sup>13</sup> was used in order to assess the level of knowledge. Then, post-intervention knowledge was evaluated in 400 participants selected randomly from these areas (participants in the pre-test were different from participants in the post-test in each areas).

The study data were entered into the SPSS statistical software (version 11.5, SPSS Inc., Chicago, IL, USA). Independent t-test was used to compare the level of awareness. Moreover, comparison of the qualitative variables between the two groups in every region was performed through chi-square and Fisher's exact tests. All P values below 0.05 were considered statistically significant.

## Results

Among the study subjects, 13 (1.6%) were male and the rest were female. In addition, 706 (88.3%) were married and 292 (36.5%) were illiterate. According to table 2, mean scores of age, marriage status, and level of education did not differ between pre-intervention and post-intervention participants ( $P > 0.05$ ). However, a difference was observed between the two groups in terms of gender ( $P = 0.003$ ). Mean scores of knowledge in pre-intervention and post-intervention groups were  $16.4 \pm 3.5$  and  $16.4 \pm 3.6$ , respectively ( $P = 0.025$ ). Mean scores of knowledge in pre-intervention and post-intervention groups did not differ between areas 1, 2, and 4. A statistically significant increase in knowledge was only observed in the area 3 group ( $P < 0.001$ ) (Table 3).

**Table 2. Demographic characteristics of participants**

Variables		Pre-intervention (mean $\pm$ SD)	Post-intervention (mean $\pm$ SD)	P
Age		39.1 $\pm$ 14.1	38.2 $\pm$ 13.8	0.388
Gender	Male	12.0 $\pm$ 3.0	1.0 $\pm$ 0.2	0.003
	Female	388.0 $\pm$ 97.0	399.0 $\pm$ 99.8	
Marital status	Single	46.0 $\pm$ 11.5	48.0 $\pm$ 12.0	0.826
	Married	354.0 $\pm$ 88.5	352.0 $\pm$ 88.0	
	Illiterate	148.0 $\pm$ 37.0	144.0 $\pm$ 36.0	
Level of education	Primary school	110.0 $\pm$ 27.5	112.0 $\pm$ 28.0	0.741
	Middle school	53.0 $\pm$ 13.2	64.0 $\pm$ 16.0	
	Diploma	64.0 $\pm$ 16.0	54.0 $\pm$ 13.5	
	Academic	25.0 $\pm$ 6.2	26.0 $\pm$ 6.5	

SD: Standard deviation

**Table 3. Comparison of the mean score of knowledge between the control and interview groups**

Type area	Educational Method	Pre-intervention (mean $\pm$ SD)	Post-intervention (mean $\pm$ SD)	P
Area 1	Based on need assessment	15.3 $\pm$ 2.8	14.9 $\pm$ 2.8	0.269
Area 2	Community-based education or local groups (CBO)	17.8 $\pm$ 3.2	16.7 $\pm$ 3.4	0.056
Area 3	Training through public classes and practical work	16.7 $\pm$ 3.2	18.4 $\pm$ 2.9	< 0.001
Area 4	Leitner method	16.2 $\pm$ 3.9	16.0 $\pm$ 3.8	0.701
Total	-	16.4 $\pm$ 3.5	16.4 $\pm$ 3.6	0.025

SD: Standard deviation

## Discussion

In the present study, training through public classes and practical work in the review of food units consumed in the previous few days, measuring anthropometric indices, informing individuals about them, and practical training on cooking was effective in increasing nutrition awareness. This study shows that chain education may not be effective. The significant difference between the two groups' mean scores of knowledge after the education by the health workers shows that the implementation of this method in the trained groups has been quite effective. The initial training was performed through various methods and the participants were asked to share the content they learned with others. However, the results showed that the chain training method is effective and encourages individuals to educate others and share their information with others. Kafeshani *et al.* investigated the mothers' knowledge status in the marginalized households of Isfahan, Iran, and showed that 20.6% of the mothers had desirable knowledge, while 79.4% had moderate and poor knowledge.<sup>13</sup> Researchers, in another study, mentioned literacy level as one of the factors effective on the level of knowledge before and after the training<sup>14</sup>. Having more knowledge can improve health-related behaviors.<sup>15</sup> Education is the foundation of all learning.<sup>16</sup> In this study, various methods were used in the training programs. Although the headlines of educational materials and intervention during the period were similar in all groups and only the training methods were different, other groups had no effects on the creation of the educational chain. Some other studies which have addressed the efficacy of various educational methods have also reported differences in the effectiveness of various methods in improving the level of knowledge.<sup>17</sup> Nevertheless, in order to design health education interventions in this province and probably all through the country, NGOs alone do not have the ability to train and health workers should be used as instructors.<sup>18,19</sup> In an

ideal condition, nutritional counseling and education should be designed based on the education level of the individual.<sup>20</sup> In addition to differences in length and method of providing the training, in many studies, educational researches are designed according to specific individuals,<sup>21,22</sup> training and consultation take place from person to person,<sup>19,23</sup> or a combination of nutritional education and behavioral counseling is presented.<sup>22</sup> Kim *et al.* demonstrated that the village health worker training program is effective in building health promotion capacity of community leaders and can be a main method for helping lower developed countries with further development<sup>24</sup>. On the other hand, educating individuals and raising their level of knowledge has been shown to be highly effective in improving their nutritional status.<sup>25,26</sup> Evidently, an individual's nutritional behavior is not only affected by his/her nutritional knowledge and can be influenced by numerous factors.<sup>27,20</sup> The present study had one limitation, i.e. the members of the second sample were not asked whether someone had transferred the trainings to them or not because the goal of the study was creating the educational chain.

## Conclusion

According to the results of this study, using health personnel with sufficient education experience is very important in NGO educational programs because they have the required level of knowledge and communication ability, and the trainees will probably trust them more.

## Conflict of Interests

Authors have no conflict of interests.

## Acknowledgements

Our appreciation goes to the Kurdistan University of Medical Sciences for their financial support, and all who participated in and helped us perform this study.



## References

1. Sarafino EP. Health Psychology: Biopsychosocial Interactions. 4<sup>th</sup> ed. New Jersey, NJ: Wiley; 2001.
2. Dostkam H, Hosseinian J, Fatehi GH. The prevalence of myocardial infarction without ST segment in patients with primary diagnosis of unstable angina in the Ardabil Bouali hospital. *J Ardabil Univ Med Sci* 2006; 6(1): 37-43. [In Persian].
3. Aggarwal BB, Harikumar KB. Potential therapeutic effects of curcumin, the anti-inflammatory agent, against neurodegenerative, cardiovascular, pulmonary, metabolic, autoimmune and neoplastic diseases. *Int J Biochem Cell Biol* 2009; 41(1): 40-59.
4. Zulli A, Lau E, Wijaya BP, Jin X, Sutarga K, Schwartz GD, et al. High dietary taurine reduces apoptosis and atherosclerosis in the left main coronary artery: association with reduced CCAAT/enhancer binding protein homologous protein and total plasma homocysteine but not lipidemia. *Hypertension* 2009; 53(6): 1017-22.
5. Sanchez-Villegas A, Delgado-Rodriguez M, Martinez-Gonzalez MA, De Irala-Estevez J. Gender, age, socio-demographic and lifestyle factors associated with major dietary patterns in the Spanish Project SUN (Seguimiento Universidad de Navarra). *Eur J Clin Nutr* 2003; 57(2): 285-92.
6. Brenn T, Njolstad I. Coronary heart disease risk factors in subjects whose brothers, sisters or husbands developed premature myocardial infarction during 12 years of follow-up. The Finnmark Study (1977-1989). *J Cardiovasc Risk* 1998; 5(5-6): 325-30.
7. Uddin SN, Begum F, Malik F, Rahman S. Coronary artery disease in young patients: clinical review and risk factor analysis. *Mymensingh Med J* 2003; 12(1): 3-7.
8. Sheikhtaheri A, Moghaddasi H. Information Managers: The Powerful Strategists of Health care Organizations in Future. *Health Inf Manage* 2008; 5(1): 83-93. [In Persian].
9. Alm-Roijer C, Stagmo M, Uden G, Erhardt L. Better knowledge improves adherence to lifestyle changes and medication in patients with coronary heart disease. *Eur J Cardiovasc Nurs* 2004; 3(4): 321-30.
10. Ferrans CE, Zerwic JJ, Wilbur JE, Larson JL. Conceptual model of health-related quality of life. *J Nurs Scholarsh* 2005; 37(4): 336-42.
11. Cottrell RR, Girvan JT, McKenzie JF. Principles and Foundations of Health Promotion and Education. San Francisco, CA: Benjamin Cummings; 2006.
12. Alavi SH, Ebrahimzadeh IS, Karimzadegan Moghadam D, Ataran M, Mehrdad R, Golestan B. An evaluation of a new rapid e-learning approach for staff in-service training at the tums. *Iranian of Higher Education* 2010; 2(3): 67-88. [In Persian].
13. Kafeshani O, Sarrafzadegan N, Nouri F, Mohammadifard N. Major dietary patterns in Iranian adolescents: Isfahan Healthy Heart Program, Iran. *ARYA Atheroscler* 2015; 11(Special 1): 61-8.
14. Heshmat R, Keshtkar A, Sheykh-ol-Eslam R, Bagheri M, Nadim A. Knowledge, Attitude and practice of households and health care staff towards nutrition and micronutrients (NUT-KAP) in provinces under the pilot study on flour fortification with Iron: study design and sampling method. *Iran J Epidemiol* 2005; 1(1): 9-16.
15. Green LW, Richard L. The need to combine health education and health promotion: the case of cardiovascular disease prevention. *Promot Educ* 1993; Spec No: 11-8.
16. Hashemyan M. Health education and relationship. Tehran, Iran: Andishe Rafeea Publication p. 39-50; 2007.
17. Turnin MC, Tauber MT, Couvaras O, Jouret B, Bolzonella C, Bourgeois O, et al. Evaluation of microcomputer nutritional teaching games in 1,876 children at school. *Diabetes Metab* 2001; 27(4 Pt 1): 459-64.
18. Buzzard IM, Asp EH, Chlebowski RT, Boyar AP, Jeffery RW, Nixon DW, et al. Diet intervention methods to reduce fat intake: nutrient and food group composition of self-selected low-fat diets. *J Am Diet Assoc* 1990; 90(1): 42-50, 53.
19. Tsorbatzoudis H. Evaluation of a planned behavior theory-based intervention programme to promote healthy eating. *Percept Mot Skills* 2005; 101(2): 587-604.
20. Tepper BJ, Choi YS, Nayga Jr RM. Understanding food choice in adult men: Influence of nutrition knowledge, food beliefs and dietary restraint. *Food Quality and Preference* 1997; 8(4): 307-17.
21. Nordevang E, Callmer E, Marmur A, Holm LE. Dietary intervention in breast cancer patients: effects on food choice. *Eur J Clin Nutr* 1992; 46(6): 387-96.
22. Shepherd R. Influences on food choice and dietary behavior. *Forum Nutr* 2005; (57): 36-43.
23. Lanza E, Schatzkin A, Ballard-Barbash R, Corle D, Clifford C, Paskett E, et al. The polyp prevention trial II: dietary intervention program and participant baseline dietary characteristics. *Cancer Epidemiol Biomarkers Prev* 1996; 5(5): 385-92.
24. Kim JM, Koh KW, Oak CH, Jung WH, Kim SH, Park DH. Assessment of village health worker training program in Tuguegarao, Philippine. *J Prev Med Public Health* 2009; 42(6): 377-85.
25. Zare H, Shojaiezhadeh D. Evaluation of education efficacy on proper nutrition knowledge and iron deficiency anemia in Mehriz city [Msc Thesis].

- Tehran, Iran: Tehran University of Medical Sciences; 2001. [In Persian].
26. Jazayeri A, Pour Moghim M, Dorosti Motlagh AR. Evaluation and Comparison of food safety and micronutrient intake in high school girls in North and South of Tehran. *Iran J Public Health* 1999; 28(4): 1-10.
27. Newman VA, Thomson CA, Rock CL, Flatt SW, Kealey S, Bardwell WA, et al. Achieving substantial changes in eating behavior among women previously treated for breast cancer-an overview of the intervention. *J Am Diet Assoc* 2005; 105(3): 382-91.